

# Delayed sedimentary response to abrupt climate change at the Paleocene-Eocene boundary, northern Spain

Duller, Robert; Armitage, J. J.; Manners, Hayley R.; Grimes, S.T.; Dunkley Jones, Tom

DOI:  
[10.1130/G45631.1](https://doi.org/10.1130/G45631.1)

License:  
Creative Commons: Attribution (CC BY)

*Document Version*  
Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*  
Duller, R, Armitage, JJ, Manners, HR, Grimes, ST & Dunkley Jones, T 2019, 'Delayed sedimentary response to abrupt climate change at the Paleocene-Eocene boundary, northern Spain', *Geology*, vol. 47, no. 2, G45631.1, pp. 159-162. <https://doi.org/10.1130/G45631.1>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Delayed sedimentary response to abrupt climate change at the Paleocene-Eocene boundary, northern Spain

Robert A. Duller<sup>1</sup>, John J. Armitage<sup>2</sup>, Hayley R. Manners<sup>3</sup>, Stephen Grimes<sup>3</sup>, Tom Dunkley Jones<sup>4</sup>

<sup>1</sup>Department of Earth, Ocean and Ecological Sciences, University of Liverpool, Liverpool L69 3BX, UK

<sup>2</sup>Institut de Physique du Globe de Paris–Sorbonne Paris Cité, Université Paris Diderot, CNRS, UMR7154, 1 rue Jussieu, 75238 Paris CEDEX 05, France

<sup>3</sup>School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth, Devon PL4 8AA, UK

<sup>4</sup>School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

## ABSTRACT

Sediment routing systems (SRSs) are a critical element of the global response to ongoing climate change. However SRS response to climate forcing is complex, fragmentary, and obscured when viewed over short, human time scales ( $10^1$ – $10^2$  yr). Over long time scales ( $>10^2$ – $10^3$  yr), the aggregated, system-wide response of SRSs to climate forcing can be gleaned with more confidence from the sedimentary record, but the nature and time scales of this aggregated response to abrupt climate change are still poorly understood. Here, we investigate the aggregated temporal response of a SRS in northern Spain to abrupt climate warming at the Paleocene-Eocene thermal maximum (PETM). Our results show that terrestrial sites in northern Spain record a temporal lag of  $16.5 \pm 7.5$  k.y. between the onset of the PETM, defined by an abrupt negative excursion in the  $\delta^{13}\text{C}$  profile, and the onset of coarse-grained deposition. Within the same SRS at a deep marine site 500 km to the west, we observe a temporal lag of  $16.5 \pm 1.5$  k.y. using an age model that is independent of that used for the terrestrial sites. These results suggest that the aggregated, system-wide response of SRSs to present-day global warming—if we take the PETM as an appropriate modern-day analogue—may persist for many millennia into the future.

## INTRODUCTION

The Paleocene-Eocene thermal maximum (PETM) is the most informative geological analogue for understanding the impact of rapid ( $<5$  k.y.) and large-magnitude ( $>4^\circ\text{C}$ ) warming on global hydrology and sediment routing systems (SRSs) (Haywood et al., 2011; Foreman et al., 2012; Carmichael et al., 2017). The PETM is associated with a large negative carbon-isotope excursion (CIE) driven by the release of  $>4000$  Pg of isotopically light carbon into the global carbon cycle (Gutjahr et al., 2017). This event is recorded in the carbon isotope ( $\delta^{13}\text{C}$ ) values of calcium carbonate minerals and organic matter from both terrestrial and marine strata (McInerney and Wing, 2011). During the PETM, changes in atmospheric and ocean-surface temperature caused dramatic and globally nonuniform change of the hydrological cycle (Bowen and Bowen, 2008; Carmichael et al., 2017). Studies of mid-latitude areas in the central United States (Foreman et al., 2012; Kraus et al., 2013; Baczynski et al., 2013) and northern Spain (Schmitz


and Pujalte, 2007; Manners et al., 2013) indicate increased seasonal precipitation during the PETM, within generally dry climate regimes. In northern Spain, there is a clear association between the CIE and an increase in the amount, and caliber, of detrital material transported to both terrestrial and marine environments (Schmitz and Pujalte, 2003, 2007; Dunkley Jones et al., 2018; Pujalte et al., 2015). A laterally extensive ( $>500$  km<sup>2</sup>), 1–7-m-thick conglomerate unit (“Claret conglomerate”) marks this change in the terrestrial environment (Schmitz and Pujalte, 2003). Similarly, in Wyoming, USA, a 30-m-thick and laterally extensive channel-belt sandstone (“boundary sandstone”) is concomitant with the PETM (Foreman, 2014); and in Colorado, USA, a similar change in sedimentary architecture is associated with the PETM (Foreman et al., 2012). The short temporal duration of PETM onset ( $<5$  k.y.) (Bowen et al., 2015; Dunkley Jones et al., 2018), and the potential for precise, high-resolution stratigraphic correlation between sections using  $\delta^{13}\text{C}$  data, enable

PETM Earth system leads and lags to be resolved at the millennial scale (Knight and Harrison, 2013). The preservation and exposure of terrestrial and deep-marine PETM sequences in the Tremp-Graus Basin of northern Spain (Schmitz and Pujalte, 2003, 2007; Domingo et al., 2009; Manners et al., 2013; Dunkley Jones et al., 2018) allows the study of system-wide response of a SRS to rapid and abrupt climate warming. We use these sequences in northern Spain (Fig. 1) to explore the aggregated, system-wide response of a PETM SRS.

## METHODS

To quantify the temporal relationship between the onset of the PETM climate perturbation and the response of the Spanish PETM SRS, we first assume that a close coupling existed between the onset of the CIE and the PETM climate perturbation. This is supported by high-resolution studies of expanded PETM sections from the New Jersey margin (USA) that find no significant lag between  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  records (Zeebe et al., 2016). Here we focus on quantifying the lag time ( $t_{\text{lag}}$ ) between the onset of the CIE and the onset of coarse-grained detrital deposition (OCD) in a study section. To do this, we first calculate a mean sedimentation rate for each section using the latest estimates of PETM CIE duration (Westerhold et al., 2018) and the measured stratigraphic thickness of the CIE at each location. To account for uncertainty in the precise identification of the CIE “core” ( $\sim 100$  k.y. duration) versus “core and recovery” ( $\sim 180$  k.y.) in the study sections, we calculate lower and upper estimates for sedimentation rate using both of these durations (Westerhold et al., 2018).

Our approach should give minimum  $t_{\text{lag}}$  values for two reasons. First, sedimentation rates in most sections increase substantially during the



**Figure 1. Site location, location map, and early Paleogene paleogeography of Spanish Pyrenees. PETM—Paleocene-Eocene thermal maximum. Section locations: CL—Claret; T—Tendrui; E—Esplugafreda; C—Campo; Z—Zumaia. Main cities: Sa—Santander; SS—San Sebastian; Vi—Vitoria; P—Pau; To—Toulouse; Tp—Trempe. Modified from Pujalte et al. (2015).**

core of the PETM (Dunkley Jones et al., 2018; John et al., 2007), and applying these higher sedimentation rate values to the onset phase will minimize calculated  $t_{\text{lag}}$  values. Second, the basal surface of the Claret conglomerate in the Spanish sections truncates underlying strata that accumulated during the onset interval of the PETM. Therefore, the preservation of a stratigraphic offset between the onset of the PETM CIE and the base of the Claret conglomerate is strong evidence for a temporal lag of SRS response. On this basis, the  $t_{\text{lag}}$  values calculated here are almost certainly minimum estimates. We note that, for our purposes, the calculation of  $t_{\text{lag}}$  is not compromised by the Sadler effect (Sadler, 1981).

To further explore the stratigraphic response of SRSs to abrupt hydrological change, we utilize a one-dimensional (1-D) sediment transport model (see the GSA Data Repository<sup>1</sup>). The model solves for the change in topography due to the transport of sediment downslope, which is a function of both local slope and surface water discharge (Armitage et al., 2016). Precipitation rate is increased from a baseline of 0.5 m yr<sup>-1</sup> following a box profile with a time to peak precipitation of 5 k.y. reflecting the time scale of PETM onset (Westerhold et al., 2018).

## RESULTS

The analysis shows that the Tendrui, Claret, and Campo sections exhibit a stratigraphic offset, but that the Esplugafreda section does not (Fig. 2). The calculated durations of the stratigraphic offset ( $t_{\text{lag}}$ ) are: Tendrui,  $t_{\text{lag}} = 13\text{--}24$  k.y.;

Claret,  $t_{\text{lag}} = 9\text{--}16$  k.y.; Esplugafreda,  $t_{\text{lag}} = 0$  k.y. The lower rates of sediment accumulation at the Esplugafreda section (0.1 m yr<sup>-1</sup>) has led to a reduced level of stratigraphic completeness (Sadler, 1981; Straub and Esposito, 2013), inhibiting proxy record preservation and precluding the identification of a resolvable  $t_{\text{lag}}$  (Foreman and Straub, 2017). We tentatively estimate the Campo  $t_{\text{lag}}$  to be 20–36 k.y.

A new age model for the deep marine segment of this SRS at Zumaia (Fig. 1), alongside a record of detrital mass accumulation rate (MAR<sub>D</sub>) (Dunkley Jones et al., 2018), allows for a unique comparison of the terrestrial and marine response to the same event. The MAR<sub>D</sub> data from the Zumaia section show an immediate response to the CIE, increasing from 1 g cm<sup>-2</sup> k.y.<sup>-1</sup> to 3 g cm<sup>-2</sup> k.y.<sup>-1</sup> over the first 5 k.y. (Fig. 2E). This elevated value is maintained for 10 k.y. before increasing abruptly to 7 g cm<sup>-2</sup> k.y.<sup>-1</sup> over a period of <4 k.y., an increase that lags behind the CIE by  $t_{\text{lag}} = 15\text{--}18$  k.y., while maximum MAR<sub>D</sub> values are attained 25–30 k.y. after the onset of the CIE (Fig. 2E). The immediate response of MAR<sub>D</sub> to the CIE is due to the greater advective length scales or transportability of finer-grained material (Ganti et al., 2014). However, the synchronicity of the arrival of coarse material at the terrestrial sites and the increase in MAR<sub>D</sub> at the deep marine site suggests that a single causative mechanism is responsible for the observed time lag. The data presented above strongly suggest that SRSs may take ~10<sup>4</sup> yr to respond to abrupt, large-magnitude climate change.


The results of the 1-D sediment transport model demonstrate that a  $t_{\text{lag}}$  can be reproduced (Fig. 3). The greater the precipitation increase over the 5 k.y. onset duration, the shorter the value of  $t_{\text{lag}}$  (Fig. 3). The truncation of individual

grain size profiles (e.g., Fig. 3B) signifies sediment bypass and non-deposition (i.e. a time gap) of the coarsest grain-size populations. We suggest that this phase of bypass is recorded in northern Spain as a 20–30-m-thick succession of fine-grained floodplain sediment that directly overlies the Claret conglomerate. The model predicts  $t_{\text{lag}}$  values of 0 k.y. at 50 km distance from the origin, 10–35 k.y. at 100 km, and 45–85 k.y. at 150 km (Fig. 3D), which is not dissimilar to the field estimates of  $t_{\text{lag}}$ . We acknowledge that these model results are approximations.

## DISCUSSION

Stratigraphic sections from the terrestrial and deep marine segments of the northern Spain PETM SRS show strong evidence for a time lag between the onset of the PETM CIE and the onset of coarse-grained deposition. The results of a 1-D sediment transport model support this. The observed  $t_{\text{lag}}$  was generated by the internal response of the SRS; so what was this mechanism? It is possible that the natural avulsion of river channels could generate a stratigraphic offset, and so  $t_{\text{lag}}$ , but the  $t_{\text{lag}}$  value would be spatially variable. The maximum value of an *avulsion-related time lag* can be approximated by a compensation time scale,  $T_c = h/r$ , (where  $h$  is channel depth and  $r$  is aggradation rate). This defines the time window over which sediment can be delivered to the majority of a river valley width (Straub et al., 2009; Wang et al., 2011), which is calculated to be  $T_c = 6.5 \pm 3.7$  k.y. for the sections in northern Spain. This value of  $T_c$  represents a maximum value, given that the time taken for a channel to visit 95% of the river valley width could be of the order of ~0.25  $T_c$  (Straub and Esposito, 2013), and channel mobility increases with increasing sediment flux (Wickert et al., 2013), and with a reduction in vegetation cover; both of which are experienced by PETM landscapes in the subtropics (Schmitz and Pujalte, 2007; Foreman, 2014). We note that  $T_c = 6.5 \pm 3.7$  k.y. also represents a minimum resolvable lag time using geochemical proxies (Foreman and Straub, 2017). Our preferred mechanism for the generation of  $t_{\text{lag}}$  is a delay in sediment transport from source area(s) to down-system locations as transport slopes adjust to the increased transport capacity of the fluvial system. This mechanism can explain a SRS-wide response. However, given the inherent complexity of landscapes and their nonlinear response to external forcing, it is possible that mountainous landscapes can act as a buffer to hydrological change (e.g., Knight and Harrison, 2013). Once this “buffering capacity” threshold is exceeded, then whole-scale landscape response, and associated increased sediment flux to areas down-system, ensues. Given this scenario, we cannot completely rule out the possibility that the time it takes to reach this threshold is manifested as a  $t_{\text{lag}}$  or is an important contributor to it. This requires further work.

<sup>1</sup>GSA Data Repository item 2019059, details of the 1-D sediment transport model and values used to calculate the duration of stratigraphic offset, is available online at [www.geosociety.org/pubs/ft2019.htm](http://www.geosociety.org/pubs/ft2019.htm), or on request from [editing@geosociety.org](mailto:editing@geosociety.org).



Given the lag times present in many of the studied PETM sections, our results suggest that it may take up to 15 k.y. for SRS to achieve a total and aggregated, system-wide response to modern abrupt climate forcing. This time scale probably represents an upper estimate given that anthropogenic carbon release rates are an order of magnitude greater than that associated with the PETM CIE (Zeebe et al., 2016). The aggregated, system-wide response time scale described here is akin to the time scale necessary for “system clearing” (Jerolmack and Paola, 2010; Foreman et al., 2012) to occur in an individual SRS following abrupt climate forcing. This temporal and spatial character of SRS


Armitage, J.J., Burgess, P.M., Hampson, G.J., and Allen, P.A., 2016, Deciphering the origin of cyclical gravel front and shoreline progradation and retrogradation in the stratigraphic record: Basin Research, v. 30, p. 15–35, <https://doi.org/10.1111/bre.12203>.

Baczynski, A.A., McInerney, F.A., Wing, S.L., Kraus, M.J., Bloch, J.I., Boyer, D.M., Secord, R., Morse, P.E., and Fricke, H.C., 2013, Chemostratigraphic implications of spatial variation in the Paleocene-Eocene Thermal Maximum carbon isotope excursion, SE Bighorn Basin, Wyoming: Geochemistry Geophysics Geosystems, v. 14, p. 4133–4152, <https://doi.org/10.1002/ggge.20265>.

Bowen, G.J., and Bowen, B.B., 2008, Mechanisms of PETM global change constrained by a new record from central Utah: Geology, v. 36, p. 379–382, <https://doi.org/10.1130/G24597A.1>.

Bowen, G.J., Maibauer, B.J., Kraus, M.J., Röhl, U., Westerhold, T., Steimke, A., Gingerich, P.D., Wing,





**Figure 3.** Model results of time delay as function of distance from source. **A:** Four “increased precipitation” scenarios.  $\Delta p$ —positive change in precipitation from initial value of  $0.5 \text{ m yr}^{-1}$ . **B,C:** Chronostratigraphic plots showing grain-size response of each precipitation scenario at two locations: at 50 km and 100 km from source. PO—Paleocene-Eocene thermal maximum (PETM) onset; OCD—onset of coarse detrital deposition; SBP—sediment bypass. Time difference between PO and OCD is time delay or lag ( $t_{\text{lag}}$ ). **D:** Model  $t_{\text{lag}}$  results (white-filled symbols) and field data results (gray-filled symbols) plotted as function of distance. T—Tendry section; CL—Claret section; C—Campo section; Z—Zumaia section;  $\text{MAR}_{\text{D(max)}}$ —Maximum detrital mass accumulation rate at Zumaia (Dunkley Jones et al., 2018).

- S.L., and Clyde, W.C., 2015, Two massive, rapid releases of carbon during the onset of the Paleocene-Eocene thermal maximum: *Nature Geoscience*, v. 8, p. 44–47, <https://doi.org/10.1038/ngeo2316>.
- Carmichael, M.J., et al., 2017, Hydrological and associated biogeochemical consequences of rapid global warming during the Paleocene-Eocene Thermal Maximum: *Global and Planetary Change*, v. 157, p. 114–138, <https://doi.org/10.1016/j.gloplacha.2017.07.014>.
- Domingo, L., López-Martínez, N., Leng, M.J., and Grimes, S.T., 2009, The Paleocene-Eocene Thermal Maximum record in the organic matter of the Claret and Tendry continental sections (South-central Pyrenees, Lleida, Spain): *Earth and Planetary Science Letters*, v. 281, p. 226–237, <https://doi.org/10.1016/j.epsl.2009.02.025>.
- Dunkley Jones, T., et al., 2018, Orbital forcing of terrestrial hydrology, weathering and carbon sequestration during the Paleocene-Eocene Thermal Maximum: *Climate of the Past*, <https://doi.org/10.5194/cp-2017-131>.
- Foreman, B.Z., 2014, Climate-driven generation of a fluvial sheet sand body at the Paleocene-Eocene boundary in north-west Wyoming (USA): *Basin Research*, v. 26, p. 225–241, <https://doi.org/10.1111/bre.12027>.
- Foreman, B.Z., and Straub, K.M., 2017, Autogenic geomorphic processes determine the resolution and fidelity of terrestrial paleoclimate records: *Science Advances*, v. 3, e1700683, <https://doi.org/10.1126/sciadv.1700683>.
- Foreman, B.Z., Heller, P.L., and Clementz, M.T., 2012, Fluvial response to abrupt global warming at the Paleocene/Eocene boundary: *Nature*, v. 491, p. 92–95, <https://doi.org/10.1038/nature11513>.
- Ganti, V., Lamb, M.P., and McElroy, B., 2014, Quantitative bounds on morphodynamics and implications for reading the sedimentary record: *Nature Communications*, v. 5, 3298, <https://doi.org/10.1038/ncomms4298>.
- Gutjahr, M., Ridgwell, A., Sexton, P.F., Anagnostou, E., Pearson, P.N., Pälike, H., Norris, R.D., Thomas, E., and Foster, G.L., 2017, Very large release of mostly volcanic carbon during the Paleocene-Eocene Thermal Maximum: *Nature*, v. 548, p. 573–577, <https://doi.org/10.1038/nature23646>.
- Haywood, A.M., Ridgwell, A., Lunt, D.J., Hill, D.J., Pound, M.J., Dowsett, H.J., Dolan, A.M., Francis, J.E., and Williams, M., 2011, Are there pre-Quaternary geological analogues for a future greenhouse warming?: *Philosophical Transactions of the Royal Society of London A*, v. 369, p. 933–956, <https://doi.org/10.1098/rsta.2010.0317>.
- Jerolmack, D.J., and Paola, C., 2010, Shredding of environmental signals by sediment transport: *Geophysical Research Letters*, v. 37, L19401, <https://doi.org/10.1029/2010GL044638>.
- John, C.M., Bohaty, S.N., Zachos, J.C., Sluijs, A., Gibbs, S., Brinkhuis, H., and Bralower, T.J., 2007, North American continental margin records of the Paleocene-Eocene thermal maximum: Implications for global carbon and hydrological cycling: *Paleoceanography*, v. 23, PA2217, <https://doi.org/10.1029/2007PA001465>.
- Knight, J., and Harrison, S., 2013, The impacts of climate change on terrestrial Earth surface systems: *Nature Climate Change*, v. 3, p. 24–29, <https://doi.org/10.1038/nclimate1660>.
- Kraus, M.J., McInerney, F.A., Wing, S.L., Secord, R., Baczynski, A.A., and Bloch, J.I., 2013, Paleohydrologic response to continental warming during the Paleocene-Eocene Thermal Maximum, Bighorn Basin, Wyoming: *Paleogeography, Palaeoclimatology, Palaeoecology*, v. 370, p. 196–208, <https://doi.org/10.1016/j.palaeo.2012.12.008>.
- Manners, H.R., et al., 2013, Magnitude and profile of organic carbon isotope records from the Paleocene-Eocene Thermal Maximum: Evidence from northern Spain: *Earth and Planetary Science Letters*, v. 376, p. 220–230, <https://doi.org/10.1016/j.epsl.2013.06.016>.
- McInerney, F.A., and Wing, S.L., 2011, The Paleocene-Eocene Thermal Maximum: A perturbation of carbon cycle, climate, and biosphere with implications for the future: *Annual Review of Earth and Planetary Sciences*, v. 39, p. 489–516, <https://doi.org/10.1146/annurev-earth-040610-133431>.
- Pujalte, V., Baceta, J.I., and Schmitz, B., 2015, A massive input of coarse-grained siliciclastics in the Pyrenean Basin during the PETM: The missing ingredient in a coeval abrupt change in hydrological regime: *Climate of the Past*, v. 11, p. 1653–1672, <https://doi.org/10.5194/cp-11-1653-2015>.
- Sadler, P.M., 1981, Sediment accumulation rates and the completeness of stratigraphic sections: *The Journal of Geology*, v. 89, p. 569–584, <https://doi.org/10.1086/628623>.
- Schmitz, B., and Pujalte, V., 2003, Sea-level, humidity, and land-erosion records across the initial Eocene thermal maximum from a continental-marine transect in northern Spain: *Geology*, v. 31, p. 689–692, <https://doi.org/10.1130/G19527.1>.
- Schmitz, B., and Pujalte, V., 2007, Abrupt increase in seasonal extreme precipitation at the Paleocene-Eocene boundary: *Geology*, v. 35, p. 215–218, <https://doi.org/10.1130/G23261A.1>.
- Straub, K.M., and Esposito, C.R., 2013, Influence of water and sediment supply on the stratigraphic record of alluvial fans and deltas: Process controls on stratigraphic completeness: *Journal of Geophysical Research: Earth Surface*, v. 118, p. 625–637, <https://doi.org/10.1002/jgrf.20061>.
- Straub, K.M., Paola, C., Mohrig, D., Wolinsky, M.A., and George, T., 2009, Compensational stacking of channelized sedimentary deposits: *Journal of Sedimentary Research*, v. 79, p. 673–688, <https://doi.org/10.2110/jsr.2009.070>.
- Wang, Y., Straub, K.M., and Hajek, E.A., 2011, Scale-dependent compensational stacking: An estimate of autogenic time scales in channelized sedimentary deposits: *Geology*, v. 39, p. 811–814, <https://doi.org/10.1130/G32068.1>.
- Westerhold, T., Röhl, U., Wilkens, R.H., Gingerich, P.D., Clyde, W.C., Wing, S.L., Bowen, G.J., and Kraus, M.J., 2018, Synchronizing early Eocene deep-sea and continental records—Cyclostratigraphic age models from the Bighorn Basin Coring Project drill cores: *Climate of the Past*, v. 14, p. 303–319, <https://doi.org/10.5194/cp-14-303-2018>.
- Wickert, A.D., Martin, J.M., Tal, M., Kim, W., Sheets, B., and Paola, C., 2013, River channel lateral mobility: Metrics, time scales, and controls: *Journal of Geophysical Research: Earth Surface*, v. 118, p. 396–412, <https://doi.org/10.1029/2012JF002386>.
- Zeebe, R.E., Ridgwell, A., and Zachos, J.C., 2016, Anthropogenic carbon release rate unprecedented during the past 66 million years: *Nature Geoscience*, v. 9, p. 325–329, <https://doi.org/10.1038/ngeo2681>.

Printed in USA